

# EXCELLENT KNOCKDOWN AND RESIDUAL PERFORMANCE


## BROAD SPECTRUM PROVIDES GREAT VALUE

With decades of proven performance, Imidan® WP Insecticide is a great option for apple, cranberry and blueberry growers.

### **EXCELLENT KNOCKDOWN & RESIDUAL PERFORMANCE**

Imidan WP Insecticide provides both quick contact activity on insect targets as well as extended residual control. Also, with minimal disruption to beneficial insects, Imidan is a great option for Integrated Pest Management.











### EFFECTS OF PH ON HALF LIFE

Insecticidal activity of Imidan is greatly reduced when the water in the spray solution is pH 6.5 or higher. It is extremely important to test water for pH levels before using for Imidan.

Acidify or buffer the tank mix solution to a pH or lower prior to adding Imidan. LI 700 from UAP is a great option.

рН	5.0	5.5	6.0	6.5	7.0	7.5	8.2
Half Life	178 hrs	92 hrs	36 hrs	14 hrs	10 hrs	2 hrs	33 min













# **CROPS**

Apples
Blueberries
Carrots
Celery
Cranberries

#### **PESTS**

Please see product label for complete list of pests controlled



1-800-960-4318 gowancanada.com



REGISTRATION NO. 29064 PEST CONTROL PRODUCTS ACT Imidan® is a registered trademark of Gowan Company L.L.C. ALWAYS READ AND FOLLOW LABEL DIRECTIONS.

IB23\_02\_Imidan\_WP\_overview\_vc001



# EXCELLENT KNOCKDOWN AND RESIDUAL PERFORMANCE

# CROPS, PESTS, RATES, REMARKS

Crops	Pests	Rates	Remarks		
Apples	Codling Moth Redbanded Leafroller Plum Curculio Apple Maggot Obliquebanded Leafroller Apple Aphid Japanese Beetle Spotted Wing Drosophila Tarnished Plant Bug	2.68 kg/ha (1.09 kg/ac)	DO NOT apply within 14 days of harvest Re-Entry Interval: 22 days for pick-your-own		
Blueberries	Blueberry Spanworm Blueberry Maggot Japanese Beetle (adults) Spotted Wing Drosophila	1.6 kg/ha (0.65 kg/ac)	DO NOT apply within 15 days of harvest.  Re-Entry Interval: 17 days for pick-your-own.  Maximum of 1 application per season.		
Carrots and Celery	Carrot Weevil	1.6 kg/ha (0.65 kg/ac)	DO NOT apply within 40 days of harvest.  Re-Entry Interval: 12 hours.  Maximum of 1 application per season.		
Cranberries	Blackheaded fireworm	1.57 kg/ha (0.64 kg/ac)	DO NOT apply within 30 days of harvest.  Re-Entry Interval: 3 days.  Maximum of 1 application per season.		

Consult label for spray volumes, additional insects and additional crop information

For more information on Imidan WP Insecticide please contact your local Gowan Canada Sales Representative, call 1-800-960-4318 or visit us online at gowancanada.com

1-800-960-4318 gowancanada.com



REGISTRATION NO. 29064 PEST CONTROL PRODUCTS ACT Imidan® is a registered trademark of Gowan Company L.L.C. ALWAYS READ AND FOLLOW LABEL DIRECTIONS. IB23\_02\_Imidan\_WP\_overview\_vc001