

BENEFITS

- Group 3 herbicide mode of action and is an excellent resistance management tool
- Reduced selection pressure for post-emergent herbicides
- Maximize yields with early weed removal

APPLICATION

Trifluralin can be tied up in straw and chaff residues, so do not apply Treflan[®] Liquid EC to fields with more than 30% crop residue coverage (70% soil). Treflan Liquid EC can be mixed with glyphosate or other burndown materials to clean up existing weeds and give residual control of broadleaf and grassy weeds.

RECROPPING

Do not seed oats, sugarbeets, timothy, canary seed, or creeping red fescue the year after using Treflan Liquid EC. Do not seed wheat if trifluralin and/or ethalfluralin were used in consecutive years.

RATES

% Organic Matter	Sandy, Sandy Loam		Loam, Silty Clay Loam, Silt Loam, Clay Loam, Silt, Silty Clay, Sandy Clay Loam, Clay	
	Spring	Fall	Spring	Fall
2 - 6% OM	1.7 L/ha (0.7 L/ac)	2.3 L/ha (0.9 L/ac)	2.3 L/ha (0.9 L/ac)	3.0 L/ha (1.2 L/ac)
6 - 15% OM	2.3 L/ha (0.9 L/ac)	3.0 L/ha (1.2 L/ac)	2.3 - 3.0 L/ha (0.9 - 1.2 L/ac)	3.0 - 3.4 L/ha (1.2 - 1.4 L/ac)**

*Fall Only

**Higher rate for heavier wild oat infestations

¹Does not include trifluralin-resistant green foxtail



CROPS

- Lentils*
- Peas
- Canola
- Mustard
- Flax*
- Dry Beans
- Soybeans
- Fababeans
- Sunflowers
- Alfalfa Establishment
- Clovers
- Birds-Foot Trefoil

WEEDS

- Green Foxtail¹
- Yellow Foxtail
- Barnyard Grass
- Japanese Brome
- Downy Brome
- Persian Darnel
- Wild Buckwheat
- Pigweed
- Lamb's-Quarters
- Cow Cackle
- Chickweed
- Purslane
- Knotweed

1-800-960-4318
gowancanada.com



Treflan[®] is a registered trademark of Gowan Company, L.L.C.
ALWAYS READ AND FOLLOW LABEL DIRECTIONS.