

BENEFITS

- Group 15 mode of action will control group 1 and 2 resistant wild oats
- Controls wild oats before they emerge
- Reduced selection pressure for post-emergent herbicides
- Maximize yields with early weed removal
- Excellent choice for base of herbicide layering program

APPLICATION TIMING

Spring and Fall are both options. Apply at the first available opportunity to make sure it gets done:

Fall: Any time after September 15th

Spring: At least 7 days before seeding

INCORPORATION

Shallow incorporation with a heavy harrow at time of application or within 24 hours will increase control, based on research studies.

BULK DENSITY AND APPLICATOR SETTINGS

629.53 kg/cu. m. (39.3 lb/cu. ft). Use the nearest number in applicator flow rate sheet. Calibration is required for accurate application rates.

RECROPPING

Do not seed oats the year following Avadex application.

SPRING & FALL RATES

22.7 KG or 451.33 KG bags available

Crop	2 - 4% OM	>4% OM	Seeding Depth
Spring or Durum Wheat, Barley, Canary Seed	14 kg/ha (12.5 lbs/ac)	17 kg/ha (15 lbs/ac)	Minimum 5 cm (1.5") (place seed 1.27 cm (1/2") below incorporation depth)
Canola, Flax, Mustard, Field Peas	17 kg/ha (15 lbs/ac)	22 kg/ha (20 lbs/ac)	As desired

Please refer to product label for complete rate details. REGISTRATION NO. 25112 PEST CONTROL PRODUCTS ACT.

For more information on Avadex[®] MicroActiv[®] Herbicide please contact your local Gowan Canada Sales Representative, call 1-800-960-4318 or visit us online at gowancanada.com



CROPS

Spring Wheat
Durum Wheat
Barley
Canary Seed
Canola
Flax
Yellow Mustard
Brown Mustard
Oriental Mustard
Sugar Beets
Field Peas

WEEDS

Wild Oats
(including Group 1
and group 2
resistant biotypes)



1-800-960-4318
gowancanada.com

Gowan[®]
CANADA

Avadex[®] is a registered trademark used under license by Gowan Company, L.L.C. MicroActiv[®] is a registered trademark of Gowan Company, L.L.C. ALWAYS READ AND FOLLOW LABEL DIRECTIONS.