

Nexter - Booklet

GROUP	21	INSECTICIDE
-------	----	-------------

NEXTER SC® Miticide/Insecticide

For control of listed mites in apples, cherries (sweet and tart), grapes, pears, strawberries and peach/nectarine orchards, listed mites as a post-harvest application on raspberries, and pear psylla in pears. Peaches and nectarines are registered in Ontario only.

COMMERCIAL

SOLUBLE CONCENTRATE

**READ THE LABEL AND BROCHURE BEFORE USING.
KEEP OUT OF REACH OF CHILDREN.**

ACTIVE INGREDIENT: Pyridaben 450 g/L

Contains 1,2-benzisothiazolin-3-one at 0.0193% as a preservative.

**REGISTRATION NO.: 33433
PEST CONTROL PRODUCTS ACT**



**WARNING POISON
EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER**

Warning, contains the allergen sulfite

NET CONTENTS: 0.5 L - Bulk

Gowan®
CANADA

Canyon Group, L.L.C.
P.O. Box 5569
Yuma, AZ 85366-5569

Product Information: 1-800-883-1844

NEXTER® is a registered trademark of Nissan Chemical Industries Limited used under license by Gowan Company, L.L.C.

GENERAL INFORMATION

This container contains **NEXTER SC Miticide/Insecticide** to control mite pests, including European red mite, apple rust mite, McDaniel spider mite, pear rust mite, twospotted spider mite, and pear psylla, when used at recommended rates.

NEXTER SC Miticide/Insecticide works primarily through contact action, so good spray coverage is necessary.

NEXTER SC Miticide/Insecticide provides excellent knockdown and residual control.

SAFETY AND HANDLING

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

1. Harmful or fatal if swallowed.
2. Harmful if inhaled. Avoid inhaling vapour or spray mist.
3. May irritate eyes and skin. Avoid contact with eyes, skin and clothing.
4. Potential skin sensitizer.
5. Wash thoroughly after handling and before eating, drinking or smoking.
6. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, during airblast application, wear protective eyewear (goggles or face shield) and chemical-resistant headgear. A chemical-resistant headgear includes chemical-resistant Sou'Wester hat, chemical-resistant rain hat or large brimmed waterproof hat and hood with sufficient neck protection. Gloves, protective eyewear and a hat are not required during application within a closed cab.
7. DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) specified in the following table. Employers should make every effort to schedule pesticide applications and worker tasks in order to avoid early entry of workers into treated areas. Under exceptional circumstances, certified pesticide applicators may enter treated areas for short-term tasks not involving hand labour if at least 4 hours have passed since application and a long-sleeved shirt, long pants, chemical-resistant footwear, socks, goggles, chemical-resistant gloves and a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides is worn. Time spent in the treated area cannot exceed 1 hour in a 24-hour period or until restricted entry interval is over.

Crop	Postapplication Activities	Restricted Entry Intervals
Grapes	Girdling, turning	54 days
	Tying, training, hand harvesting, leaf pulling by hand	30 days
	Mechanical harvesting ¹	25 days
	All other activities	24 hours
All other crops	All activities	24 hours

¹Mechanical harvesting assumes no contact with treated foliage or crop.

8. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
9. Clean spray equipment thoroughly after use.

10. Do not contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.
11. Do not use in residential areas. Residential areas are defined as sites where bystanders including children may be potentially exposed during or after spraying. This includes around homes, school, parks, playgrounds, playing fields, public buildings or any other areas where the general public including children could be exposed.
12. Hazardous to humans and domestic animals. Keep out of reach of children and pets.
13. This product is only to be used by individuals holding an appropriate pesticide applicator certificate or license recognized by the provincial/territorial pesticide regulatory agency where the pesticide application is to occur.
14. This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not apply or allow **NEXTER SC Miticide/Insecticide** to drift to crops or weeds on which bees are actively foraging. Do not allow spray to drift towards beehives.
15. Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca.

ENVIRONMENTAL PRECAUTIONS:

TOXIC to bees. Bees may be exposed through direct spray, spray drift, and residues on leaves, pollen and nectar in flowering crops and weeds. Minimize spray drift to reduce harmful effects on bees in habitats close to the application site. Avoid applications when bees are foraging in the treatment area in ground cover containing blooming weeds. To further minimize exposure to pollinators, refer to the complete guidance "Protecting Pollinators during Pesticide Spraying – Best Management Practices" on the Health Canada website (www.healthcanada.gc.ca/pollinators). Follow crop specific directions for application timing.

For applications on crops that are highly attractive to pollinators (apples, raspberries, cherries, nectarines, peaches, and pears), and crops where managed bees are present: DO NOT apply during the crop blooming period.

For applications to grapes and strawberries: Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging.

TOXIC to beneficial arthropods. Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland.

TOXIC to birds.

TOXIC to small wild mammals.

TOXIC to aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.

TOXIC to non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

In case of emergency involving a major spill, leak or fire call Chemtrec® at 1-800-424-9300. For medical emergencies call 1-888-478-0798

TOXICOLOGICAL INFORMATION

The patient should be treated symptomatically.

STORAGE

1. Store in original tightly closed container.
2. Do not ship or store near food, feed, seed and fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.
4. To prevent contamination store this product away from food or feed.

DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

If this is a recyclable container:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.

2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

If this is a returnable container:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

If this is a refillable container:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

DIRECTIONS FOR USE

Please refer to and retain this booklet for complete directions.

MIXING

1. Fill the spray tank one-half full of water and start agitation.
2. Add NEXTER SC Miticide/Insecticide to the tank.
3. Continue agitation while filling tank to the desired spray volume.
4. Dilute **NEXTER SC Miticide/Insecticide** in enough water to provide thorough spray coverage. If foaming occurs, an anti-foaming agent may be used.

It is recommended that a small number of plants be treated prior to application to the entire planting. Nexter SC miticide/insecticide may leave visible residue when used at high rates.

TANK MIXING

Physical incompatibility, reduced pest control or crop injury may result from mixing **NEXTER SC Miticide/Insecticide** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers. Gowan Company does not offer any recommendations for tank mixing **NEXTER SC Miticide/Insecticide** with any other products.

RESTRICTIONS AND LIMITATIONS

Do not apply this product through any type of irrigation system.

DO NOT APPLY THIS PRODUCT BY AIR.

Do not apply directly to aquatic systems. Aquatic systems include all permanent lentic (standing) and lotic (flowing) water bodies. This product is toxic to aquatic invertebrates, fish and aquatic plants.

TIMING

NEXTER SC Miticide/Insecticide should be applied when populations are building and before serious crop injury occurs. Consult provincial guidelines and local extension experts for treatment threshold and advice. Mites cause the most economic damage one week to one month following bloom. **NEXTER SC Miticide/Insecticide** provides excellent knockdown and residual control. A good performance evaluation can be made 7-10 days after treatment.

CROPS

Apples

Cherries (sweet and tart)

Peaches/Nectarines (Ontario only)

Strawberries

Grapes

Apply to grapes when immature motile stages are present for best control.

Pears

Apply when pear psylla or mite numbers reach significant levels. Second application to pears for mites and pear psylla can be made 30 days after the first application.

Raspberries

Apply to raspberries after harvest when mites appear.

The use of **NEXTER SC Miticide/Insecticide** is compatible with the goal of preservation of predatory mites. This product is easiest on predatory mites when applied late spring to early summer before numbers of beneficial mites have peaked. **NEXTER SC Miticide/Insecticide** is harmful to certain other beneficials (i.e. beneficial insects).

RECOMMENDED APPLICATION RATES

Table 1. NEXTER SC Miticide/Insecticide Application Rate Table

Mites Controlled	Rate*
European Red Mite (<i>Panonychus ulmi</i>) Apple Rust Mite (<i>Aculus schlechtendali</i>) Pear Rust Mite (<i>Epirimerus pyri</i>)	500 ml/ha (225 g ai/ha)
Twospotted Spider Mite (<i>Tetranychus urticae</i>) McDaniel Spider Mite	500 ml/ha** (225 g ai/ha) or

<i>(Tetranychus mcdanieli)</i>	1000 ml/ha *** (450 g ai/ha)
Pear Psylla <i>(Cacopsylla pyricola)</i>	1000 ml/ha – 1200 ml/ha (450 g ai/ha – 540 g ai/ha)

* Based on water volume (3000 L/ha).

** Good control will be achieved at or below the mite action threshold.

*** For faster knockdown and control above mite action thresholds late season.

NOTE: NEXTER SC Miticide/Insecticide controls nymph and larva stage of two-spotted spider and McDaniel mites and all motile stages of European red, apple rust mite and pear rust mite. This product is not an ovicide.

If product is applied as a concentrate spray, do not apply at rates less than 330 ml/ha (148.5 g ai/ha) for European red, apple rust mite, pear rust mite, two-spotted and McDaniel spider mite or 650 ml/ha (292.5 g ai/ha) for two-spotted and McDaniel spider mite above action threshold level.

Do not use less than 1000 L/ha water volume. For foliar application using ground application equipment only.

Crop Recommendation Table

Crop	Application to Harvest Interval (Days)	Maximum Number of Applications/ Year	Minimum Application Interval (Days)	Maximum Product Used (L/ha)/ Year on the Same Orchard
Apples	25	2	30	2.0
Cherries (sweet and tart)*	7	1	NA	1.0
Grapes	25	1	NA	1.0
Peaches/Nectarines (Ontario only)	14	1	NA	1.0
Pears	25	2	30	2.4
Raspberries: Post-harvest application only	N/A	2	30	2.0
Strawberries	10	2	15	2.0

*For optimal control, apply product in sufficient water for good leaf coverage (3000 L/ha).

DO NOT apply by hand-held mistblower or hand-held fogger.

DO NOT apply by aerial application equipment.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 8 km/h at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of

Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Air-induction nozzles must be used for the ground application of this product.

Airblast application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. **DO NOT** apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

Buffer zones:

Spot treatments using hand-held equipment **DO NOT** require a buffer zone.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrub-lands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Buffer Zones (metres) Required for the Protection of:				
			Freshwater Habitat of Depths:		Estuarine/Marine Habitat of Depths:		Terrestrial habitat
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Groundboom	Strawberry, raspberry		20	2	1	1	1
Airblast	Pear, apple, raspberry	Late growth stage	65	40	30	20	1
	Cherry, peach, nectarine	Late growth stage	60	35	30	20	0

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The buffer zones for airblast application of this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pest Management Regulatory Agency web site. Buffer zones for groundboom sprayer application **CANNOT** be modified using the Buffer Zone Calculator.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, please note that **NEXTER SC Miticide/Insecticide** contains a Group 21 insecticide/acaricide. Any insect/mite population may contain individuals naturally

resistant to **NEXTER SC Miticide/Insecticide** and other Group 21 insecticides/acaricides. The resistant individuals may dominate the insect/mite population if this group of insecticides/acaricides is used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of **NEXTER SC Miticide/Insecticide** or other Group 21 insecticides/acaricides with different groups that control the same pests in a field.
- Use tank mixtures with insecticides/acaricides from a different group when such use is permitted.
- Insecticide/acaricide use should be based on an IPM program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance, contact Gowan Company at 1-800-883-1844.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Nexter – base panel for booklet – permanently attached to foil bag

GROUP

21

INSECTICIDE

NEXTER® SC Miticide/Insecticide

For control of listed mites in apples, cherries (sweet and tart), grapes, pears, strawberries and peach/nectarine orchards, listed mites as a post-harvest application on raspberries, and pear psylla in pears. Peaches and nectarines are registered in Ontario only.

**COMMERCIAL
SOLUBLE CONCENTRATE**

**READ THE LABEL AND BROCHURE BEFORE USING.
KEEP OUT OF REACH OF CHILDREN.**

ACTIVE INGREDIENT: Pyridaben 450 g/L

Contains 1,2-benzisothiazolin-3-one at 0.0193% as a preservative.

**REGISTRATION NO.: 33433
PEST CONTROL PRODUCTS ACT**



**WARNING POISON
EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER**

Warning, contains the allergen sulfite

NET CONTENTS: 0.5L - Bulk

Gowan®
CANADA

Gowan Company, L.L.C.
P.O. Box 5569
Yuma, AZ 85366-5569

Product Information: 1-800-883-1844

NEXTER® is a registered trademark of Nissan Chemical Industries Limited used under license by Gowan Company, L.L.C.

NEXTER® SC Miticide/Insecticide

For control of listed mites in apples, cherries (sweet and tart), grapes, pears, strawberries and peach/nectarine orchards, listed mites as a post-harvest application on raspberries, and pear psylla in pears. Peaches and nectarines are registered in Ontario only.

**COMMERCIAL
SOLUBLE CONCENTRATE**

**READ THE LABEL AND BROCHURE BEFORE USING.
KEEP OUT OF REACH OF CHILDREN.**

ACTIVE INGREDIENT: Pyridaben 450 g/L

Contains 1,2-benzisothiazolin-3-one at 0.0193% as a preservative.

**REGISTRATION NO.: 33433
PEST CONTROL PRODUCTS ACT**



**WARNING POISON
EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER**

Warning, contains the allergen sulfite

NET CONTENTS: 0.5 L - Bulk

Gowan®
CANADA

Gowan Company, L.L.C.
P.O. Box 5569
Yuma, AZ 85366-5569

Product Information: 1-800-883-1844

NEXTER® is a registered trademark of Nissan Chemical Industries Limited used under license by Gowan Company, L.L.C.

GENERAL INFORMATION

This container contains **NEXTER SC Miticide/Insecticide** to control mite pests, including European red mite, apple rust mite, McDaniel spider mite, pear rust mite and twospotted spider mite, and pear psylla, when used at recommended rates

SAFETY AND HANDLING

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

1. Harmful or fatal if swallowed.
2. Harmful if inhaled. Avoid inhaling vapour or spray mist.
3. May irritate eyes and skin. Avoid contact with eyes, skin and clothing.
4. Potential skin sensitizer.
5. Wash thoroughly after handling and before eating, drinking or smoking.
6. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, during airblast application, wear protective eyewear (goggles or face shield) and chemical-resistant headgear. A chemical-resistant headgear includes chemical-resistant Sou'Wester hat, chemical-resistant rain hat or large brimmed waterproof hat and hood with sufficient neck protection. Gloves, protective eyewear and a hat are not required during application within a closed cab.
7. DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) specified in the following table. Employers should make every effort to schedule pesticide applications and worker tasks in order to avoid early entry of workers into treated areas. Under exceptional circumstances, certified pesticide applicators may enter treated areas for short-term tasks not involving hand labour if at least 4 hours have passed since application and a long-sleeved shirt, long pants, chemical-resistant footwear, socks, goggles, chemical-resistant gloves and a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides is worn. Time spent in the treated area cannot exceed 1 hour in a 24-hour period or until restricted entry interval is over.

Crop	Postapplication Activities	Restricted Entry Intervals
Grapes	Girdling, turning	54 days
	Tying, training, hand harvesting, leaf pulling by hand	30 days
	Mechanical harvesting ¹	25 days
	All other activities	24 hours
All other crops	All activities	24 hours

¹Mechanical harvesting assumes no contact with treated foliage or crop.

8. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
9. Clean spray equipment thoroughly after use.
10. Do not contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.

11. Do not use in residential areas. Residential areas are defined as sites where bystanders including children may be potentially exposed during or after spraying. This includes around homes, school, parks, playgrounds, playing fields, public buildings or any other areas where the general public including children could be exposed.
12. Hazardous to humans and domestic animals. Keep out of reach of children and pets.
13. This product is only to be used by individuals holding an appropriate pesticide applicator certificate or license recognized by the provincial/territorial pesticide regulatory agency where the pesticide application is to occur.
14. This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not apply or allow **NEXTER SC Miticide/Insecticide** to drift to crops or weeds on which bees are actively foraging. Do not allow spray to drift towards beehives.
15. Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca.

ENVIRONMENTAL PRECAUTIONS:

TOXIC to bees. Bees may be exposed through direct spray, spray drift, and residues on leaves, pollen and nectar in flowering crops and weeds. Minimize spray drift to reduce harmful effects on bees in habitats close to the application site. Avoid applications when bees are foraging in the treatment area in ground cover containing blooming weeds. To further minimize exposure to pollinators, refer to the complete guidance "Protecting Pollinators during Pesticide Spraying – Best Management Practices" on the Health Canada website (www.healthcanada.gc.ca/pollinators). Follow crop specific directions for application timing.

For applications on crops that are highly attractive to pollinators (apples, raspberries, cherries, nectarines, peaches, and pears), and crops where managed bees are present: DO NOT apply during the crop blooming period.

For applications to grapes and strawberries: Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging.

TOXIC to beneficial arthropods. Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland.

TOXIC to birds.

TOXIC to small wild mammals.

TOXIC to aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.

TOXIC to non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

In case of emergency involving a major spill, leak or fire call Chemtrec® at 1-800-424-9300. For medical emergencies call 1-888-478-0798

TOXICOLOGICAL INFORMATION

The patient should be treated symptomatically.

STORAGE

1. Store in original tightly closed container.
2. Do not ship or store near food, feed, seed and fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.
4. To prevent contamination store this product away from food or feed.

DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

If this is a recyclable container:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

If this is a returnable container:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

If this is a refillable container:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.